

บรรณานุกรม

มหาวิทยาลัยราชภัฏวชิรเวศน์บุรีรัมย์

บรรณานุกรม

- กล้าณรงค์ ศรีรอด และ เกื้อกุล ปิยะจอมขวัญ. (2546). เทคโนโลยีของแป้ง. กรุงเทพฯ : สำนักพิมพ์มหาวิทยาลัยเกษตรศาสตร์.
- เกตุการ ดาจันทร์, ธวัชชัย ศุภวิทิตพัฒนา, ปิยวรรณ ศุภวิทิตพัฒนา, อุทัยวรรณ จัตตวง, และ เอกชัย ชูเกียรติโรจน์. (2555). การผลิตมกกล้าเชื้อและผงปรุงรสจากถั่วเหลืองหมักเพื่อการยกระดับอาหารพื้นบ้านสู่การแข่งขันในตลาดธุรกิจ. รายงานการวิจัยฉบับสมบูรณ์. พิษณุโลก : มหาวิทยาลัยราชภัฏพิบูลสงคราม.
- ชุตินุช สุจริต. (2544). การเตรียมมกกล้าเชื้อผงสำหรับการผลิตปลาต้ม. รายงานการวิจัยการเตรียมมกกล้าเชื้อผงสำหรับการผลิตปลาต้ม. กระทรวงศึกษาธิการ : สถาบันเทคโนโลยีราชมงคล.
- เพิ่มพงษ์ ศรีประเสริฐ. (2524). การผลิตและการเก็บเชื้อแบคทีเรียแลคติกที่ใช้เป็นอาหารเสริมสุกรในรูปเชื้อผง. วิทยานิพนธ์ปริญญาโท. วิทยาศาสตร์มหาบัณฑิตสาขาวิชาจุลชีววิทยา มหาวิทยาลัยเกษตรศาสตร์.
- วรรณทิตา ลาภศิริ และ ศรีเวียง ทิพกานนท์. (2539). การแยกเชื้อแบคทีเรียแลคติกจากหน่อไม้เปรี้ยวเพื่อใช้เป็นกกล้าเชื้อ. รายงานการวิจัย กรุงเทพมหานคร : คณะอุตสาหกรรมเกษตร มหาวิทยาลัยเทคโนโลยีพระจอมเกล้าพระนครเหนือ.
- สมชาย ประภาวัต. (2535). เทคโนโลยีในการทำแป้งถั่วเหลือง. กรุงเทพฯ : สถาบันค้นคว้าและพัฒนาผลิตภัณฑ์อาหาร มหาวิทยาลัยเกษตรศาสตร์.
- อรอนงค์ นัยวิกุล. (2550). ข้าวสาลี : วิทยาศาสตร์และเทคโนโลยี. กรุงเทพมหานคร : มหาวิทยาลัยเกษตรศาสตร์.
- AOAC. (2000). **Official methods of analysis of AOAC International.** (17th ed). Gaithersburg, MD : AOAC International.
- Campbell-Platt, G. (1980). African locust bean (*Parkia* species) and its West African fermented food products, dawadawa. **Ecology of Food and Nutrition.** 9: 123-132.
- Chukeatirote, E., Chainun, C., Siengsubchart, A., Moukamnerd, C., Chantawannakul, P., Lumyong, S., Boontim, N., & Thakang, P. (2006). Microbiological and biochemical changes in *Thua Nao* fermentation. **Research Journal of Microbiology.** 1 : 38-44.

- Crisafulli A., Altavilla D., & Marini H. (2005). Effects of the phytoestrogen genistein on cardiovascular risk factors in post-menopausal women. **Menopause**. 12 : 186-192.
- Dajanta, K. (2010). **Production of high nutritional fermented soybean (*Thua Nao*) by *Bacillus subtilis***. Dissertation Food Science and Technology, Faculty of Agro-Industry, Chiang Mai University.
- Dajanta, K., Apichartsrangkoon, A., & Chukeatirote, E. (2011a). Antioxidant properties and total phenolics of *Thua Nao* (a Thai fermented soybean) as affected by *Bacillus*-fermentation. **Journal of Microbial and Biochemical and Technology**. 3(4) : 056-059.
- _____. (2011b). Volatile profiles of *thua nao*, a Thai fermented soy products. **Food Chemistry**. 125 : 464-470.
- _____. (2011c). Free amino acid profiles of *Thua Nao*, a Thai fermented soybean. **Food Chemistry**. 125 : 342-247.
- Dajanta, K., Chukeatirote, E., Apichartsrangkoon, A., & Frazier, R.A. (2009). Enhanced aglycone production of fermented soybean products by *Bacillus* species. **Acta Biologica Szegediensis**. 53 : 93-98.
- Dakwa, S., Sakyi-Dawson, E., Diako, C., Annan, N. T., & Amoa-Awua, W. K. (2005). Effect of boiling and roasting on the fermentation of soybeans into dawadawa (soy-dawadawa). **International Journal of Food Microbiology**. 104 : 69-82.
- Dike, E. N., & Odunfa, S. A. (2003). Microbiological and biochemical evaluation of a fermented soyabean product (soy-dadawadwa). **Journal of Food Science and Technology**. 40 : 606-610.
- Eden, J. (1998). Phytoestrogens and the menopause. **Balliere's Clinical Endocrinology and Metabolism**. 12 : 581-587.
- Farzana, K., Shah, S.N.H., Butt, F.B. & Awan, S.B. (2005). Biosynthesis of bacitracin solid-state fermentation by *Bacillus licheniformis* using defatted oil seed cakes as substrate. **Pakistan Journal of Pharmaceutical Sciences**. 18(1) : 55-57.
- Georgetti, S. R., Vicentini, F. T. M. C., Yokoyama, C. Y., Borin, M. F., & Spadaro, A. C. C. (2009). Enhanced *in vitro* and *in vivo* antioxidant activity and mobilization of free phenolic compounds of soybean flour fermented with different β -glucosidase-producing fungi. **Journal of Applied Microbiology**. 106 : 459-466.

- Gotoh, T., Yamada, K., Yin, H., Ito, A., Kataoka, T., & Dohi, K. (1998). Chemoprevention of *N*-Nitroso-*N*-methylurea-induced rat mammary carcinogenesis by soy foods or biochanin A. **Japanese Journal Cancer Research**. 89 : 137–142.
- Griffith, A. P., & Collisn, M. W. (2001). Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid chromatography-mass spectrometry. **Journal of Chromatography**. 913 : 397-413.
- Hermansen K., Sondergaard M., Hoie L., Carstensen M., & Brock B. (2001). Beneficial effects of a soy-based dietary supplement on lipid levels and cardiovascular risk markers in type 2 diabetic subjects. **Diabetes Care**. 24 : 228-233.
- Hosoi, T., & Kiuchi, K. (2003). **Natto-A Food Made by Fermenting Cooked Soybeans with *Bacillus subtilis* (natto)**. In E. R. Farnworth (Ed.), Handbook of Fermented Functional Foods (pp. 267-290). Boca Raton : CRC Press.
- Ishimi, Y., Yoshida, M., Wakimoto, S., Wu, J., Chiba, H., Wang, X., Takeda, K., & Miyaura, C. (2002). Genistein, a soybean isoflavone, affects bone marrow lymphopoiesis and prevents bone loss in castrated male mice. **Bone**. 31 : 180-185.
- Jideani, I. A. O., & Okeke, C. R. (1991). Comparative-study of microorganisms and sensory attributes of condiments from the fermentation of different seeds. **Plant foods for Human Nutrition**. 41 : 27-34.
- Jung, K.-O., Park, S.-Y., & Park, K.-Y. (2006). Basic nutritional investigation: longer aging time increases the anticancer and antimetastatic properties of *doenjang*. **Nutrition**. 22 : 539–545.
- Kim, N. Y., Song, E. J., Kwon, D. Y., Kim, H. P., & Heo, M. Y. (2008). Antioxidant and antigenotoxic activities of Korean fermented soybean. **Food and Chemical Toxicology**. 46 : 1184-1189.
- Kim, S. L., Chi, H. Y., Kim, J. T., Lee, Y. H., Park, N. K., Son, J. R., & Kim, S. J. (2006). Isoflavone content and its relationship with other seed quality traits of soybean cultivars collected in South Korea. **Korean Journal of Crop Science**. 51 : 81–88.

- Kiuchi, K., and Watanabe, S. (2004). **Industrialization of Japanese natto**. In K. H. Steinkraus (Ed.), *Industrialization of Indigenous Fermented Foods*. (pp. 193-246). (2nd ed). Marcel Dekker : New York.
- Kunioka M. & Goto A. (1994). Biosynthesis of poly (γ -glutamic acid) from L-glutamic acid, citric acid, and ammonium sulfate in *Bacillus subtilis* IFO3335. **Applied Microbiology and Biotechnology**. 40 : 867-872.
- Kwak, C. S., Lee, M. S., & Park, S. C. (2007). Higher antioxidant properties of Chungkukjang, a fermented soybean paste, may be due to increased aglycone and malonylglycoside isoflavone during fermentation. **Nutrition Research**. 27 : 719-727.
- Lee, M. Y., Park, S.-Y., Jung, K.-O., Park, K.-Y., & Kim, S. D. (2005). Quality and functional characteristics of Chungkukjang prepared with various *Bacillus* sp. isolated from traditional Chungkukjang. **Journal of Food Science**. 70 : M191-M196.
- Leejeerajumnean, A. (2003). Thua nao: alkali fermented soybean from *Bacillus subtilis*. **Silpakorn University International Journal**. 3 : 277-292.
- Liu, D., Zhen, W., Yang, Z., Carter, J.D., Si, H., & Reynolds, K.A. (2006). Genistein acutely stimulates insulin secretion in pancreatic β -cells through a cAMP-dependent protein kinase pathway. **Diabetes**. 55 : 1043-1050.
- Makela S. I., Pylkkänen L. H., Santti R. S., & Adlercreutz H. (1995). Dietary soybean may be antiestrogenic in male mice. **Journal of Nutrition**. 125 : 437-445.
- Moktan, M.R., Norbu, L., Nirola, H., Dukpa, K., Rai, T.B., & Dorji, R. (2008). Ecological and social aspects of transhumant herding in Bhutan. **Mountain Research and Development**. 28(1) : 41-48.
- Morabito N, Crisafulli A., & Vergara C. (2002). Effects of genestein and hormone-replacement therapy on bone loss in early postmenopausal women: A randomized double blind placebo-controlled study. **Journal of Bone Mineral Research**. 17 : 1904-1912.
- Nikkuni, S., Karki, T.B., Vilku, K. S., Suzuki, T., Shindoh, K., Suzuki, C., & Okada, N. (1995). Mineral in Nepal and amino acid contents of Kinema, a fermented soybean food prepared in Nepal. **Food Science and Technology International**. 1 : 107-111.

- Nout, M. J. R., Bakshi, D., & Sarkar, P. K. (1998). Microbiological safety of kinema, a fermented soya bean food. **Food Control**. 9 : 357-362.
- Ogawa Y, Yamaguchi F, Yuasa K, & Tahara Y. (1997). Efficient production of – polyglutamic acid by *Bacillus subtilis* (natto) in jar fermenters. **Bioscience, Biotechnology, and Biochemistry**. 61 : 1684-1687.
- Ohta T. (1986) **Natto**. In Legume-Based Fermented Foods, eds. N.R. Reddy, M.D. Pierson, and D.K. Salunkhe, pp. 85-95. Boca Raton, FL: CRC Press.
- Omafuvbe, B. O. (2008). Effect of temperature on biochemical changes induced by *Bacillus subtilis* (SDA3) during starter culture fermentation of soybean into condiment (soy-Daddawa). **American Journal of Food Technology**. 3 : 33-41.
- Omafuvbe, B. O., Abiose, S. H., & Shonukan, O. O. (2002). Fermentation of soybean (*Glycine max*) for soy-daddawa production by starter cultures of *Bacillus*. **Food Microbiology**. 19 : 561-566.
- Park, K.-Y., Jung, K.-O., Rhee, S.-H., & Choi, Y. H. (2003). Antimutagenic effects of *doenjang* (Korean fermented soypaste) and its active compounds. **Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis**. 523–524 : 43–53.
- Potter, S. M., Baum, J. A., Teng, H., Stillman, R. J., Shay, N. F., & Erdman, J. W. Jr. (1998). Soy protein and isoflavones: their effects on blood lipids and bone density in postmenopausal women. **The American Journal of Clinical Nutrition**. 68 : 1375S-1379S.
- Sarkar, M., Snibbe, S. S., Tversky, O. J., & Reiss, S. P. (1993). **Stretching the rubber sheet: A metaphor for viewing large layouts on small screens**. In Proceedings of UIST. 93 : 81-91.
- Sarkar, P. K., Jones, L. J., Craven, G. S., Somerset, S. M., & Palmer, C. (1997). Amino acid profiles of kinema, a soybean-fermented food. **Food Chemistry**. 59 : 69-75.
- Sarkar, P. K., & Tamang, J. P. (1995). Changes in the microbial profile and proximate composition during natural and controlled fermentation of soybeans to produce kinema. **Food Microbiology**. 12 : 317-325.

- Setchell K.D.R. (1998). Phytoestrogens: the biochemistry, physiology, and implications for human health of isoflavones. **American Journal of Clinical Nutrition**. 68 : 1333s-1346s.
- Smith, A.K., & Circle, S.J. (1972). **Soybean : Chemistry and Technology**. AVI Publishing Co. Inc.
- Steinkraus, K.H. (1991). **African alkaline Fermented Foods and their relationship to similar foods in other parts of the world**. Westby, A. & Reilly, J.A. eds. pp 87-92 Traditional African Foods-Quality and Nutrition Stockholm, Sweden. IFS.
- Sundhagul, M., Smanmathuroj, P., & Bhodacharoen, W. (1972). Thua-Nao: A fermented soybean food of northern Thailand. I. Traditional processing method. **Thai Journal of Agricultural Science**. 5 : 43-56.
- Taira, H. (1992). Quality and its variation on soybeans in Japan. **Journal of the Japanese Society for Food Science and Technology**. 39: 122-133.
- Taira, H., Tanaka, H., Saito, M. & Saito, M. (1990). Effect of cultivar, seed size and crop year on total and free sugar contents of domestic soybean. **Journal of the Japanese Society for Food Science and Technology**. 37: 203-213.
- Tamang, J. P., & Nikkuni, S. (1996). Selection of starter cultures for the production of kinema, a fermented soybean food of the Himalaya. **World Journal of Microbiology and Biotechnology**. 12 : 629-635.
- Tamang, J. P., Sarkar, P. K., & Hesselstine, C. W. (1988). Traditional fermented foods and beverages of Darjeeling and Sikkim-a review. **Journal of the Science of Food and Agriculture**. 4 : 375-385.
- Teng, D.F., Lin, C.S., & Hsieh, P.C. (2004). Fermented Whole Soybean and Soybean Paste. In **Handbook of Food and Beverage Fermentation Technology**. Marcel Dekker, Inc. United States of America.
- Tseng, Y. H., Lee, Y. L., Li, R. C., & Mau, J. L. (2005). Non-volatile flavor components of *Ganoderma tsugae*. **Food Chemistry**. 90 : 409-415.
- Tsuji, K. & Tsuji, E. (1986). Effect of natto-feeding on cholesterol level of rats. **Japanese Journal of Nutrition**. 44 : 41-44.
- Visessanguan. W., S. Benjakul. W. Potachareon, A. Panya, & S. Riebroy. (2005). Accelerated proteolysis of soy proteins during fermentation of Thua-nao inoculated with *Bacillus subtilis*. **Journal of Food Biochemistry**. 29(4): 349-366.

- Wang, D., Wang, L.-J., Zhu, F.-X., Zhu, J.-Y, Chen, X.D., Zou, L., Saito, M., & Li, L.-T. (2008). *In vitro* and *in vivo* studies on the antioxidant activities of the aqueous extracts of Douchi (a traditional Chinese salt-fermented soybean food). **Food Chemistry**. 107 : 1421-1428.
- Wei, Q.-K., Chen, T.-R., & Chen, J.-T. (2008). Use of *Bacillus subtilis* to enrich isoflavone aglycones in fermented natto. **Journal of the Science of Food and Agriculture**. 88 : 1007-1011.
- Yoshida, T. (1988). **Marine algae of Japan**. Tokyo, Japan : Uchida Rokakuho.
- Zhao, S., Hu, N., Huang, J., Liang, Y. and Zhao, B. (2008). High-yield spore production from *Bacillus licheniformis* by solid state fermentation. **Biotechnology Letters**. 30: 295-297.